

## Left ventricular unloading before reperfusion reduces endothelin-1 release and calcium overload in porcine myocardial infarction

Sophie Tamareille, PhD,<sup>a</sup> Hela Achour, MD,<sup>a</sup> James Amirian, BA,<sup>a</sup> Patricia Felli, BS,<sup>a</sup> Roger J. Bick, PhD,<sup>b</sup> Brian Poindexter, MS,<sup>b</sup> Yong J. Geng, MD, PhD,<sup>c</sup> William H. Barry, MD,<sup>c</sup> and Richard W. Smalling, MD, PhD, FACC<sup>a</sup>

**Objectives:** The aim of this study was to test the hypothesis that after an acute myocardial infarction, endothelin-1 release with subsequent calcium overload is a mediator of myocardial reperfusion injury, which can be inhibited, in part, by left ventricular unloading immediately before reperfusion. We recently have reported that left ventricular unloading before reperfusion reduces infarct size after acute myocardial infarction. However, the biologic mechanisms of infarct salvage in unloaded hearts subjected to ischemia/reperfusion remain undefined.

**Methods:** Twelve pigs were subjected to 1 hour of left anterior descending coronary artery occlusion followed by 4 hours of reperfusion. A left ventricular assist device was initiated 15 minutes before reperfusion and maintained during reperfusion (assist device group, n = 6). A control group (n = 6) was subjected to reperfusion alone. Infarct size, endothelin-1 plasma levels, intracellular calcium levels, and apoptosis were analyzed in both groups.

**Results:** At reperfusion, left ventricular unloading significantly decreased left ventricular end-diastolic and end-systolic pressures. Infarct size, expressed as a percentage of zone at risk, was also significantly reduced by 54% in the group with the left ventricular assist device compared with controls. Support with a left ventricular assist device reduced endothelin-1 release from the heart at 15 minutes, 30 minutes, and 1 hour of reperfusion. Myocardial release of endothelin-1 was significantly correlated with infarct size at 15 minutes of reperfusion ( $r = 0.79$ ;  $P = .008$ ). Left ventricular unloading caused a significant reduction of calcium overload and of the percentage of apoptotic cells in the ischemic region.

**Conclusion:** Our findings suggest that endothelin-1 release and calcium overload are important mediators of reperfusion injury and that they can be significantly reduced by left ventricular unloading before coronary artery reperfusion during myocardial infarction.

Prompt institution of reperfusion after coronary artery occlusion limits myocardial infarct size.<sup>1-3</sup> However, reperfusion after more than 1 hour of ischemia, a common clinical scenario, may result in significant “reperfusion injury,” which diminishes the extent of myocardial salvage.<sup>1</sup> Pharmacologic treatment alone, such as the early use of beta-blockers, nitroglycerin, and angiotensin-converting enzyme inhibitors, has not been effective in the prevention of reperfusion injury.<sup>4-6</sup> Increased circulating plasma levels and myocardial tissue content of endothelin-1

From the Division of Cardiovascular Medicine<sup>a</sup> and Department of Pathology,<sup>b</sup> University of Texas at Houston Medical School, Houston, Tex, and University of Utah Health Science Center,<sup>c</sup> Salt Lake City, Utah.

This study was supported in part by a grant from A-Med Systems, Inc, West Sacramento, Calif, and generous contributions from numerous grateful patients.

Received for publication Sept 10, 2007; revisions received Jan 4, 2008; accepted for publication Jan 29, 2008.

Address for reprints: Richard W. Smalling, MD, PhD, FACC, Department of Internal Medicine, Division of Cardiovascular Medicine, University of Texas Houston Medical School, 6431 Fannin, MSB 1.246, Houston, TX 77030 (E-mail: richard.w.smalling@uth.tmc.edu).

J Thorac Cardiovasc Surg 2008;136:343-51  
0022-5223/\$34.00

Copyright © 2008 by The American Association for Thoracic Surgery

doi:10.1016/j.jtcvs.2008.01.021

**Abbreviations and Acronyms**

Ca <sup>2+</sup>	= calcium
CS	= coronary sinus
ET-1	= endothelin-1
INF/LV	= infarct weight/left ventricular weight ratio
IV	= intravenous
LAD	= left anterior descending coronary artery
LV	= left ventricle/ventricular
LVAD	= left ventricular assist device
MI	= myocardial infarction
TUNEL	= terminal deoxynucleotidyltransferase-mediated dUTP-biotin nick end labeling
ZR/LV	= zone at risk/left ventricular weight ratio

(ET-1), a 21-amino acid peptide with a powerful vasoconstrictor activity, have been detected during ischemia/reperfusion, and a possible role for this peptide in the pathophysiology of ischemia/reperfusion injury has been postulated.<sup>7,8</sup>

Mechanical unloading of the myocardium reduces left ventricular (LV) pressure work and myocardial oxygen consumption.<sup>9-12</sup> We<sup>13</sup> have previously shown that LV unloading during myocardial infarction (MI) limits reperfusion injury as well as infarct size independent of its effects on myocardial blood flow. We<sup>13</sup> have also demonstrated that LV unloading initiated just before reperfusion produced a dramatic reduction in calcium (Ca<sup>2+</sup>) overload and contraction band necrosis by electron microscopy in the ischemic region compared with reperfusion alone. Others have shown in isolated myocytes subjected to simulated ischemia that ET-1 promotes Ca<sup>2+</sup> flux into the cells, which is associated with cell death.<sup>14</sup> We therefore tested the hypothesis that LV unloading just before reperfusion reduces ET-1 release on reperfusion, and we proposed that the reduced ET-1 levels would then reduce Ca<sup>2+</sup> flux into ischemic myocytes, reducing fatal Ca<sup>2+</sup> overload and consequently reducing infarct size.

**Materials and Methods****Animal Model**

We used 12 Yorkshire pigs of either sex weighing 50 ± 5 kg. The study protocol was approved by the Animal Welfare Committee of the University of Texas at Houston Medical School, and all experiments were performed according to the Committee's guidelines.

**Surgical Procedure**

Pigs were sedated with tiletamine HCl and zolazepam HCl (Telazol IM; Wyeth, Madison, NJ, 2.0 mg/kg, to effect) and intubated. General anesthesia was maintained with isoflurane (0.5–2.5 vol%). An 8F sheath was placed in the right carotid artery and a catheter was advanced to the ascending aorta for monitoring aortic pressure and for blood sampling for regional myocardial blood flow and ET-1 sampling. A 6F right multipurpose catheter was placed into the coronary sinus (CS) under fluoroscopic guidance. The A-Med left ventricular

assist device (LVAD) (A-Med Systems, Inc, West Sacramento, Calif), a miniaturized circulatory assist device, was inserted into the LV via femoral access as previously described.<sup>13</sup> In control animals, the introducer sheath was inserted in the femoral artery but the pumping was not initiated. The heart was exposed via a sternotomy. A ligature was placed around the left anterior descending coronary artery (LAD) at a position from which the distal third of the artery would be occluded by tightening the ligature. A Doppler flow probe was placed around the artery just proximal to the snare for monitoring coronary blood flow. A catheter was placed in the LV via an apical stab wound for measurement of LV pressure. Catheters were placed in the left atrium for measurement of atrial pressure and for radioactive microsphere injection. All animals received acetylsalicylic acid (325 mg intravenously [IV]) 30 minutes before ischemia, and sodium heparin 5000 IU IV bolus followed by an IV drip to maintain an activated clotting time of greater than 200 seconds until the end of the experiment. An IV lidocaine drip of 1 mg/min was started and maintained throughout the occlusion period for 2 hours into reperfusion or until the animal's condition had stabilized. In addition, bretylium tosylate was infused before the occlusion with an IV drip rate of 5 mg · kg<sup>-1</sup> · h<sup>-1</sup> maintained throughout the procedure. Hemodynamic and functional measurements were recorded on a Gould chart recorder (ThermoSpectra Corp, San Diego, Calif) and analyzed as previously described.<sup>13</sup> After baseline measurements, the LAD was occluded and the occlusion was maintained for 1 hour followed by reperfusion for 4 hours. Flow cessation in the LAD was documented by the Doppler flow signal. At the experiment conclusion, the animals were heparinized (4000 IU IV), fully anesthetized with intravenous pentobarbital, and humanely killed with supersaturated potassium chloride. The heart was quickly excised from the chest and washed with tap water.

**Experimental Protocol**

Baseline measurements included hemodynamic values (arterial blood pressure, left atrial pressure, and LV pressure), coronary blood flow, and arterial and CS ET-1 sampling. Hemodynamic values and flow data were continuously recorded. Blood samples for measuring plasma ET-1 were collected simultaneously from the CS and the descending thoracic aorta at baseline, 45 minutes of ischemia, reperfusion, and 15 and 30 minutes and 1, 2, and 4 hours of reperfusion. Tissue biopsy specimens for Ca<sup>2+</sup> measurements were taken at baseline, 45 minutes of ischemia, and at 10 minutes, 30 minutes, and 4 hours of reperfusion. Group 1 (n = 6) served as a control group and was not subjected to LV assistance. In group 2 (n = 6), LV unloading was started 15 minutes before reperfusion at maximal rotational speed and maintained for the remainder of the experiment.

**Plasma ET-1 Content**

ET-1 concentration was determined by the radioimmunoassay technique with the isoform-specific rabbit antibodies against synthetic ET-1 (Peninsula Laboratories, Inc, San Carlos, Calif). Blood samples were collected simultaneously from the CS and the aorta and processed according to the manufacturer's instructions.

**Assessment of Intracellular Ca<sup>2+</sup> in Cardiac Tissue**

Myocardial tissue biopsy specimens, 13-mm long, were obtained from the ischemic and control regions with an 18-gauge needle (Bio-Pince; Amedic AB, Sollentuna, Sweden) for Ca<sup>2+</sup> measurements.

Intracellular Ca<sup>2+</sup> content was determined with the Ca<sup>2+</sup>-sensitive fluorescent probe, fluo-3 AM (Molecular Probes, Eugene, Ore), using a fluorescence deconvolution microscope (Applied Precision Delta Vision, Issaquah, Wash), incorporating an Olympus IX70 microscope (Olympus America, Melville, NY), as previously described.<sup>15</sup> A combination of probes was used to determine cell types (myocytes, vascular endothelial cells, and fibroblasts). 4,6-Diamino-2-phenylindole (DAPI) was used to identify nuclei. Smooth muscle actin, cardiac muscle actin, and cardiac myosin were probed with secondary antibody tagged with DIPYromethene BOron Difluoride (BODIPY) or Texas Red dye. Smooth muscle actin identified vascular elements, whereas actin and myosin patterns and absence of intercalated disks distinguished myocytes from fibroblasts.<sup>15</sup>

**Infarct Size Determination**

Infarct size quantification by triphenyl tetrazolium chloride staining was measured according to the method of Fishbein and associates.<sup>16</sup> After excision of the heart, the aortic root was perfused with Evans blue dye (1%) while the LAD distal to the occlusion was perfused with triphenyl tetrazolium chloride at equal pressures. The heart was then sliced. The blue-stained control region, the red-stained risk region, and the tan infarct were measured by planimetry and quantified in all slices as described previously.<sup>9</sup>

**Regional Myocardial Blood Flow**

Regional myocardial blood flow was analyzed with the microsphere technique at baseline, 25 minutes of ischemia, and 5 minutes of reperfusion and was expressed as milliliters per gram per minute.<sup>17</sup> Flows were measured in the endocardial and epicardial areas as well as transmurally.

**TUNEL Assay**

Terminal deoxynucleotidyltransferase-mediated dUTP-biotin nick end labeling (TUNEL)-positive cardiomyocytes were detected in

paraffin-embedded LV sections by the protocol described in the DeadEnd Colorimetric TUNEL System (Promega Corporation, Madison, Wis). Positive and negative control sections were included. Microscopic evaluations were performed by a BX40 inverted microscope (Olympus America).

**Statistical Analysis**

All results are expressed as mean ± SEM. For multiple comparison procedures, including hemodynamic and coronary flow data, infarct size, ET-1 levels, and Ca<sup>2+</sup> measurements, analysis of variance and the Newman-Keuls multiple-range tests were performed (Statistica; StatSoft, Inc, Tulsa, Okla). Analysis of covariance was performed to compare infarct sizes, with ET-1 levels or collateral blood flow used as a covariate.

**Results**

**Hemodynamic Changes**

Hemodynamic changes during ischemia and reperfusion are shown in Table 1. At baseline and during LAD occlusion and reperfusion, there was no significant difference in heart rate, mean arterial pressure, or rate–pressure product between control and unloaded animals. Initiation of the pump resulted in LV unloading, as indicated by significant decreases in end-diastolic and systolic pressures.

**Myocardial Blood Flow**

Myocardial blood flow, expressed as millimeters per gram per minute, was measured in the area of coronary occlusion as well as in the contralateral area (Table 2). There was a significant decrease in myocardial blood flow with ischemia. However, there was no significant difference in absolute transmural blood flow between controls and treated animals

**TABLE 1. Hemodynamic parameters in control and unloaded animals**

	LAD occlusion				Reperfusion							
	Baseline	10 min pump off	35 min pump off	45 min pump on	10 min pump on	30 min pump on	1 h pump on	2 h pump on	3 h pump on	4 h pump on	4 h pump off	
HR (beats/min)												
Control	113 ± 4	103 ± 3	101 ± 6	94 ± 3	95 ± 2	94 ± 3	92 ± 3	94 ± 5	95 ± 7	99 ± 8	93 ± 8	
LVAD	100 ± 8	89 ± 9	94 ± 7	95 ± 9	87 ± 6	85 ± 6	81 ± 7	90 ± 8	98 ± 8	102 ± 12	86 ± 13	
MAP (mm Hg)												
Control	93 ± 8	90 ± 8	84 ± 7	88 ± 9	89 ± 10	82 ± 9	80 ± 8	64 ± 7	57 ± 5	45 ± 5	40 ± 4	
LVAD	89 ± 11	92 ± 13	91 ± 12	84 ± 11	73 ± 10	66 ± 12	67 ± 12	68 ± 15	66 ± 10	67 ± 13	54 ± 15	
LVSP (mm Hg)												
Control	109 ± 9	107 ± 9	101 ± 8	106 ± 10	109 ± 11	101 ± 10	99 ± 9	84 ± 7	79 ± 5	66 ± 6	62 ± 4	
LVAD	102 ± 10	105 ± 14	105 ± 12	71 ± 16	42 ± 9*	49 ± 12*	40 ± 12*	55 ± 12	55 ± 9*	41 ± 5*	71 ± 16	
LVEDP (mm Hg)												
Control	7 ± 1	8 ± 1	9 ± 1	10 ± 2	9 ± 1	9 ± 1	9 ± 1	7 ± 1	8 ± 1	7 ± 1	6 ± 1	
LVAD	6 ± 0	8 ± 1	8 ± 1	5 ± 2	4 ± 1*	3 ± 1*	4 ± 1*	3 ± 1*	5 ± 1	3 ± 1*	8 ± 2	
RPP												
Control	121 ± 8	109 ± 9	103 ± 9	100 ± 10	104 ± 12	95 ± 11	92 ± 9	79 ± 9	76 ± 8	65 ± 8	58 ± 8	
LVAD	104 ± 16	97 ± 18	99 ± 15	82 ± 12	66 ± 10*	63 ± 13	59 ± 14	67 ± 14	73 ± 8	67 ± 6	66 ± 16	

Values are expressed as means ± SEM of 6 experiments. HR, Heart rate; LVAD, left ventricular assist device; MAP, mean arterial pressure; LVSP, left ventricular systolic pressure; LVEDP, left ventricular end-diastolic pressure; RPP, heart rate–pressure product. \*P < .05 vs control.

**TABLE 2. Regional myocardial blood flow (transmural) in control and unloaded animals**


	Absolute flows (mL · g <sup>-1</sup> · min <sup>-1</sup> )			P
	Control	LVAD		
		Off	On	
Baseline				
Control region	0.85 ± 0.09	0.87 ± 0.09	—	.86
Ischemic region	0.62 ± 0.06	0.70 ± 0.06	—	.36
LAD occlusion				
Control region	0.70 ± 0.10	1.12 ± 0.28	—	.12
Ischemic region	0.02 ± 0.01	0.05 ± 0.02	—	.16
Reperfusion (5 min)				
Control region	0.74 ± 0.13	—	0.57 ± 0.10	.34
Ischemic region	2.50 ± 0.36	—	2.75 ± 0.45	.67

Values are means ± SEM of 6 experiments. LAD, Left anterior descending coronary artery.


in the ischemic or control regions at baseline, at 25 minutes of occlusion, and at 5 minutes of reperfusion. Endocardial and epicardial flows were also similar between the groups (data not shown).

**Infarct Size**

No differences were noted in the zone at risk/LV weight ratio (ZR/LV) ( $P = .12$ ), or infarct weight/total LV weight ratio (INF/LV) ( $P = .09$ ) between the groups (Figure 1). There was a significant (54%) reduction of infarct size when expressed as a percent of zone at risk (INF/ZR) in the supported group compared with controls ( $24.10\% \pm 6.45\%$  vs  $51.59\% \pm 6.64\%$ ;  $P = .014$ ). When the transmural collateral blood flow to the ischemic region was used as a covariate, infarct



**Figure 1. Bar graph showing infarct size (INF), expressed as a percent of zone at risk (ZR), and zone at risk and infarct zone as a percentage of the LV weight for the LVAD and control groups. Data are expressed as mean ± SEM of 6 independent experiments in each group.**




**Figure 2. Relationship between infarct size, expressed as a percentage of zone at risk, and collateral blood flow during coronary occlusion in the transmural ischemic zone. By analysis of covariance, a significant reduction in infarct size occurred with unloading, independent of collateral blood flow ( $P = .025$ ).**


size (as a percentage of area at risk) was significantly smaller in the LVAD group than in the control group ( $P = .025$ ). A significant reduction in infarct size occurred independent of collateral blood flow (Figure 2).

**Plasma ET-1 Levels**

The transcardiac release of ET-1 was significantly lower in the unloaded animals than in the controls at 15 minutes, 30 minutes, and 1 hour of myocardial reperfusion (Figure 3). To assess the relationship between myocardial ET-1 release and infarct size, independent of any other variables, we analyzed this relationship for the total group of tested animals using an analysis of covariance. Figure 4 shows the correlation between myocardial release of ET-1 at 15 minutes of



**Figure 3. Myocardial ET-1 plasma levels (coronary sinus minus aorta [Cs - Ao]) expressed in picograms per 100 µL in the control and unloaded animals (n = 6 pigs in each group).**



**Figure 4. Correlation between myocardial release of ET-1 at 15 minutes of reperfusion and final infarct size. There was a significantly positive correlation between myocardial release of ET-1 and infarct size with an  $r = 0.79$ , and a  $P = .008$ . CS – AO, Coronary sinus minus aorta.**

reperfusion and final infarct size. There was a significantly positive correlation between myocardial ET-1 release and infarct size, regardless of animals' treatment, indicating that myocardial ET-1 release was an independent predictor of infarct size at 15 minutes of reperfusion ( $r = 0.79$ ;  $P = .008$ ) (Figure. 4).

**Ischemic Zone Apoptosis**



LV unloading reduced the number of apoptotic cells in ischemic zones. Of 400 nuclei counted in the central ischemic region of the unloaded animals, 15.15% ± 6.4% displayed TUNEL-positive staining. In contrast, for the same number of nuclei, 56.69% ± 5.17% displayed TUNEL-positive staining in the ischemic region of the control animals ( $P = .01$ ) (Figure 5).

**Myocardial Ca<sup>2+</sup> Content**

When we compared ischemic and nonischemic ratios of fluo-3 fluorescence (Figure 6, top and bottom), there was no significant change in intracellular Ca<sup>2+</sup> content during reperfusion in the unloaded group. However, in the control group, a significant rise in the ischemic/nonischemic intracellular Ca<sup>2+</sup> ratio was seen in the reperfusion period compared with the occlusion period ( $4.86 ± 1.20$  vs  $1.18 ± 0.28$ ;  $P = .016$ ).

**Discussion**

We found that the transcardiac release of ET-1 was significantly lower in the unloaded animals as opposed to the untreated animals during reperfusion. This study, for the first time, demonstrated normalization of plasma ET-1 levels in the ischemic myocardium by hemodynamic unloading of the LV.



**Figure 5. Effect of LV unloading on ischemia/reperfusion-induced apoptosis in the heart. Apoptotic cardiomyocyte nuclei appear brown-stained, whereas TUNEL-negative nuclei appear blue with hematoxylin and eosin. Heavy staining of numerous TUNEL-positive cardiomyocytes was observed in the central ischemic region of the LV of the sham-operated pig after 1 hour of LAD ischemia and 4 hours of reperfusion. In contrast, few TUNEL-positive cells were detected in the LV ischemic region of the unloaded animal (arrow) (n = 3 to 4 counts for 6 hearts in each group).**

In the management of acute MI, the primary focus has been to achieve early reperfusion. This intervention can reduce infarct size, but it is successful only when reperfusion is achieved early after occlusion.<sup>18</sup> When reperfusion is delayed, infarct size increases and reperfusion injury ensues. Several groups have shown beneficial effects of mechanical unloading during acute MI.<sup>11,19,20</sup> Complete mechanical unloading of the LV with cardiopulmonary bypass, LV venting, and cardioplegia before reperfusion has led to a 77%

CSP




Figure 6. Effect of LV unloading on myocardial  $Ca^{2+}$  content. *Top*, Fluorescence deconvolution images of LV sections (ischemic region) probed with fluo-3 fluorescence. *Bottom*, Ischemic/non-ischemic ratios of fluo-3 fluorescence during occlusion and reperfusion period in control and unloaded animals. Data are expressed as mean  $\pm$  SEM of 12 to 15 slices of 4 hearts in each group.

GSP

reduction of infarct size.<sup>11</sup> This technique, however, is not applicable as a primary approach for MI owing to the significant time delays associated with initiating total cardiopulmonary bypass in patients having an MI. A more practical approach would be percutaneous unloading of the LV. Although it has been reported that chronic LVAD unloading results in regression of hypertrophy and improvement of

myocyte function and LV geometry, the cellular and molecular mechanisms responsible for these beneficial effects, particularly in acute ischemia, remain undefined. Preliminary experience in our laboratory in canine ischemia/reperfusion injury has demonstrated that LV unloading, intravenous administration of liposomal prostaglandin E<sub>1</sub>, intracoronary heparin distal to the coronary occlusion, and, to a certain

extent, intracoronary adenosine just before reperfusion provide significant infarct salvage, improvement in ischemic collateral blood flow, and reduction in contraction band necrosis paralleling a reduction in the ischemic region end-diastolic wall thickness during early reperfusion.<sup>9,13</sup> Some investigators suggested that the exaggerated flow of  $\text{Ca}^{2+}$  into the myocytes results in what is called “contraction band necrosis,”<sup>21</sup> but the causes of enhanced  $\text{Ca}^{2+}$  loading have not been defined.

ET-1 was first described as a vasoactive peptide<sup>22</sup> released from vascular endothelium. ET-1 is released from ischemic myocardium during reperfusion and may contribute to ischemia/reperfusion injury in the intact heart.<sup>23</sup> ET-1 has been shown to be involved in the overload-induced release of  $\text{Ca}^{2+}$ .<sup>24</sup> ET-1 can influence myocyte intracellular  $\text{Ca}^{2+}$  concentration by several mechanisms: (1) It stimulates  $\text{Ca}^{2+}$  release from internal stores<sup>25</sup>; (2) it enhances L-type  $\text{Ca}^{2+}$  current through the activation of protein kinase  $\text{C}^{26}$ ; (3) it activates the  $\text{Na}^+-\text{H}^+$  exchanger<sup>27</sup>; and (4) it activates the outward  $\text{I}_{\text{NaCa}}$  (ie, reverse mode  $\text{Na}^+-\text{Ca}^{2+}$  exchanger, in which  $\text{Ca}^{2+}$  influx is induced).<sup>28</sup>  $\text{Ca}^{2+}$  entry via reverse mode  $\text{Na}^+-\text{Ca}^{2+}$  exchanger is a major cause of ischemia/reperfusion injury.<sup>29</sup> It has been shown that activation of ET receptors on myocytes may trigger phospholipase C activity, resulting in hydrolysis of inositol phosphates and the subsequent release of intracellular  $\text{Ca}^{2+}$ . This leads to an increased susceptibility of myocardium to ischemia and reperfusion injury. ET-1–induced rise in intracellular  $\text{Ca}^{2+}$  concentration then causes cell hypercontracture and cell death.<sup>30</sup> In this study, we found that LV unloading, just before reperfusion, resulted in a significant inhibition in the rise of  $\text{Ca}^{2+}$  induced by reperfusion.

Similar to our findings, Morawietz and associates<sup>31</sup> showed up-regulation of the ET-A receptors in patients with heart failure that was normalized by LV unloading. It is likely that ET-1 is synthesized de novo and produced locally in the ischemic heart.<sup>32</sup> Endothelial cells, vascular smooth muscle cells, and cardiomyocytes all are potential sources of ET-1 production. A study using in situ hybridization and immunohistochemical analyses showed that cardiomyocytes are the main site of ET-1 synthesis and production in porcine ischemic hearts.<sup>33</sup> An increased myocardial expression of ET-1 messenger RNA has been observed in porcine cardiomyocytes subjected to ischemia, suggesting stimulated cardiac production during MI.<sup>33</sup> Galiuto and co-workers<sup>34</sup> showed, in a similar canine model, that LU 135252, an ET<sub>A</sub> receptor blocker, when given intravenously at the time of reperfusion, significantly limited the increase in end-diastolic wall thickness, which was also associated with a reduction in myocardial necrosis. We have now demonstrated a significant positive correlation of the transcardiac ET-1 gradient in the ischemic region with myocardial infarct size at 15 minutes of reperfusion, indicating that ET-1 is released across the ischemic heart and promotes LV damage

via ET-1 receptors in these animals. Support with the LVAD, just before reperfusion, resulted in a significant reduction in final infarct size independent of regional myocardial blood flow. Moreover, these findings indicate that the transcardiac release of ET-1 is a significant predictor of myocardial infarct size independent of regional myocardial blood flow and the degree of LV unloading. These results are consistent with the findings of Meyns and colleagues,<sup>35</sup> who demonstrated that in a sheep ischemia/reperfusion model known to be devoid of preformed coronary collaterals, “partial LV support” with a modest decrease in preload resulted in a significant reduction in infarct size without any measured increase in myocardial perfusion in the ischemic area. In isolated cultured cardiomyocytes, ET-1 expression is induced by stretch.<sup>36</sup> During acute MI, sarcomere stretching by diastolic Laplace overloading may lead to an increase in synthesis and release of ET-1 locally in the myocardium. A significant correlation between diastolic wall stress and ET-1 expression in the remote area of infarcted rat heart has been documented,<sup>37</sup> suggesting that stretching of the myocardium during diastole contributes significantly to the induction of ET-1 in acute MI. Also, the mechanical support provided by LVAD unloading has been shown to induce a significant reduction in LV wall stress/stretch.<sup>38</sup>

Another key finding of the present study was that LV unloading just before reperfusion resulted in a reduction in apoptotic cells in the ischemic and infarcted areas when compared with untreated animals. Programmed cell death plays a central role in the context of diverse heart diseases of ischemic and nonischemic origin. Recent evidence indicates that simulated ischemia followed by simulated reperfusion results in cardiomyocyte death with typical apoptotic features.<sup>39</sup> It is now widely established that after ischemia/reperfusion, both necrosis and apoptosis may contribute independently to infarct size.<sup>40</sup> After MI, the occurrence of apoptosis has been demonstrated in the infarct area and in the border zone of the infarction, whereas necrotic cells are mainly found in the central zone of the infarct.<sup>41</sup> Myocyte apoptosis appears to be the prevailing form of cell death after MI. Conflicting results exist concerning the role of ET-1 in cardiomyocyte apoptosis. Oie and associates<sup>42</sup> showed that bosentan, a mixed ET<sub>A</sub>/ET<sub>B</sub> receptor antagonist, attenuated cardiomyocyte cell loss through apoptosis in the viable peri-infarcted area (area at risk) after MI in rats. On the other hand, ET-1 has been reported to be an antiapoptotic factor in vascular smooth muscle cells and endothelial cells.<sup>43</sup> In addition, administration of ET-1 before ischemia has been shown to result in a preconditioning-like cardioprotective effect.<sup>44</sup> ET-1 appears to promote apoptosis only under extreme conditions and instead exerts antiapoptotic influence over a range of physiologic concentrations.

Evidence now indicates that  $\text{Ca}^{2+}$  overload may promote cell death.<sup>45</sup> In vitro experiments have shown that addition of intracellular  $\text{Ca}^{2+}$  buffering agents or extracellular  $\text{Ca}^{2+}$

chelators significantly inhibit caspase activation, DNA fragmentation, and apoptotic cell death.<sup>45</sup> In contrast, agents that directly mobilize Ca<sup>2+</sup>, such as Ca<sup>2+</sup> ionophores, have been shown to trigger apoptosis in diverse cell types.<sup>45</sup> Cell death caused by Ca<sup>2+</sup> overload has been implicated in reperfusion injury of myocardium after ischemia.<sup>46</sup> Moreover, it has been demonstrated that addition of Ca<sup>2+</sup> blockers abrogates apoptosis in cardiac myocytes subjected to chemical hypoxia, further supporting the claim that Ca<sup>2+</sup> influx plays a significant role in apoptotic cell death.<sup>47</sup> Our finding indicates some potential of LV unloading as an intervention to the vicious cycle of distention-induced proapoptotic phenotype shifts.

### Study Limitations

We chose to study the effect of unloading using a porcine model of myocardial ischemia/reperfusion to avoid a collateral flow artifact. In a species like the dog, naturally occurring collateral flow can significantly contribute to cell viability in both the permanently occluded and reperfused heart. However, the degree of collateral circulation in patients varies enormously and most often is not predictable. The pig, on the other hand, has virtually no epicardial collateral connections and only sparse endocardial connections.<sup>48</sup> Inasmuch as it has no pre-existing coronary collateral vessels, its infarctions have sharply defined borders, and coronary anatomy varies little between pigs.<sup>49</sup>

The question that remains to be resolved in this study is whether ET release accounts for the injury or whether the release of ET-1 adds to the injury, resulting from the altered hemodynamics that accompanies ischemia/reperfusion injury. The use of ET antagonism would help to address this question and could establish the cause–effect relationship between ET release and infarction and determine whether reducing ET release per se is a protective maneuver or whether it requires the physical unloading of the LVAD. A straightforward approach would be the use of an ET receptor antagonist, which would possibly abolish any difference between control and unloaded animals.

### Conclusion

Mechanical unloading of the LV, just before reperfusion, reduces infarct size in a porcine ischemia/reperfusion model. A possible mechanism for this effect is a reduction of myocardial release of ET-1 by LV unloading before reperfusion, which, in turn, reduces Ca<sup>2+</sup> overload into ischemic myocytes as well as apoptosis in the ischemic region. Taken together, our findings suggest that ET-1 release and Ca<sup>2+</sup> overload are important mediators of reperfusion injury and that they can be significantly reduced by LV unloading before coronary artery reperfusion during MI. The potential for developing new therapeutic strategies aimed at preventing reperfusion injury in patients subjected to myocardial revascularization in the setting of acute MI is significant.

### References

1. Reimer KA, Lowe JE, Rasmussen MM, Jennings RB. The wavefront phenomenon of ischemic cell death. 1. Myocardial infarct size vs duration of coronary occlusion in dogs. *Circulation*. 1977;56:786-94.
2. Christian TF, Schwartz RS, Gibbons RJ. Determinants of infarct size in reperfusion therapy for acute myocardial infarction. *Circulation*. 1992;86:81-90.
3. Kloner RA, Ellis SG, Lange R, Braunwald E. Studies of experimental coronary artery reperfusion. Effects on infarct size, myocardial function, biochemistry, ultrastructure and microvascular damage. *Circulation*. 1983;68(2 Pt 2):18-15.
4. Galcera-Tomas J, Castillo-Soria FJ, Villegas-Garcia MM, Florenciano-Sánchez R, Sánchez-Villanueva JG, de La Rosa JA, et al. Effects of early use of atenolol or captopril on infarct size and ventricular volume: a double-blind comparison in patients with anterior acute myocardial infarction. *Circulation*. 2001;103:813-9.
5. Jugdutt BI. Intravenous nitroglycerin unloading in acute myocardial infarction. *Am J Cardiol*. 1991;68:52D-63D.
6. Pfeffer MA, Greaves SC, Arnold JM, Glynn RJ, LaMotte FS, Lee RT, et al. Early versus delayed angiotensin-converting enzyme inhibition therapy in acute myocardial infarction. The healing and early afterload reducing therapy trial. *Circulation*. 1997;95:2643-51.
7. Velasco CE, Turner M, Inagami T, Atkinson JB, Virmani R, Jackson EK, et al. Reperfusion enhances the local release of endothelin after regional myocardial ischemia. *Am Heart J*. 1994;128:441-51.
8. Stewart DJ, Kubac G, Costello KB, Cernacek P. Increased plasma endothelin-1 in the early hours of acute myocardial infarction. *J Am Coll Cardiol*. 1991;18:38-43.
9. Smalling RW, Cassidy DB, Barrett R, Lachterman B, Felli P, Amirian J. Improved regional myocardial blood flow, left ventricular unloading, and infarct salvage using an axial-flow, transvalvular left ventricular assist device. A comparison with intra-aortic balloon counterpulsation and reperfusion alone in a canine infarction model. *Circulation*. 1992;85:1152-9.
10. Merhige ME, Smalling RW, Cassidy D, Barrett R, Wise G, Short J, et al. Effect of the hemopump left ventricular assist device on regional myocardial perfusion and function. Reduction of ischemia during coronary occlusion. *Circulation*. 1989;80(5 Pt 2):III158-66.
11. Allen BS, Okamoto F, Buckberg GD, Bugyi H, Leaf J. Reperfusion conditions: critical importance of total ventricular decompression during regional reperfusion. *J Thorac Cardiovasc Surg*. 1986;92:605-12.
12. Dennis C, Hall DP, Moreno JR, Senning A. Reduction of the oxygen utilization of the heart by left heart bypass. *Circ Res*. 1962;10:298-305.
13. Achour H, Boccalandro F, Felli P, Amirian J, Uthman M, Buja M, et al. Mechanical left ventricular unloading prior to reperfusion reduces infarct size in a canine infarction model. *Catheter Cardiovasc Interv*. 2005;64:182-92.
14. Rodriguez-Larrain JM, Sharma P, Sugishita K, Nonaka A, Yauney KD, Barry WH. Effects of endothelin-1 on cation concentrations and viability of ventricular myocytes during ATP depletion: assessment by flow cytometry. *Anal Pharmacol*. 2001;2:1-8.
15. Shan K, Bick RJ, Poindexter BJ, Nagueh SF, Shimoni S, Verani MS, et al. Altered adrenergic receptor density in myocardial hibernation in humans: a possible mechanism of depressed myocardial function. *Circulation*. 2000;102:2599-606.
16. Fishbein MC, Meerbaum S, Rit J, Lando U, Kanmatsuse K, Mercier JC, et al. Early phase acute myocardial infarct size quantification: validation of the triphenyl tetrazolium chloride tissue enzyme staining technique. *Am Heart J*. 1981;101:593-600.
17. Heymann MA, Payne BD, Hoffman JI, Rudolph AM. Blood flow measurements with radionuclide-labeled particles. *Prog Cardiovasc Dis*. 1977;20:55-79.
18. Vandeplassche G, Hermans C, Van Haecke J, Wouters L, Borgers M, Flameng W. Evaluation of factors influencing myocardial infarct size in unconscious dogs. *Cardiovasc Res*. 1991;25:844-54.
19. Axelrod HI, Galloway AC, Murphy MS, Laschinger JC, Grossi EA, Baumann FG, et al. A comparison of methods for limiting myocardial infarct expansion during acute reperfusion—primary role of unloading. *Circulation*. 1987;76(5 Pt 2):V28-32.

20. Catinella FP, Cunningham JN Jr, Glassman E, Laschinger JC, Baumann FG, Spencer FC. Left atrium-to-femoral artery bypass: effectiveness in reduction of acute experimental myocardial infarction. *J Thorac Cardiovasc Surg.* 1983;86:887-96.
21. Cooley DA, Reul GJ, Wukasch DC. Ischemic contracture of the heart: "stone heart." *Am J Cardiol.* 1972;29:575-7.
22. Yanagisawa M, Kurihara H, Kimura S, Tomobe Y, Kobayashi M, Mitsui Y, et al. A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature.* 1988;332:411-5.
23. Monti LD, Allibardi S, Piatti PM, Valsecchi G, Costa S, Pozza G, et al. Triglycerides impair postischemic recovery in isolated hearts: roles of endothelin-1 and trimetazidine. *Am J Physiol Heart Circ Physiol.* 2001;281:H1122-30.
24. Sugden PH. An overview of endothelin signaling in the cardiac myocyte. *J Mol Cell Cardiol.* 2003;35:871-86.
25. Yang HT, Sakurai K, Sugawara H, Watanabe T, Norota I, Endoh M. Role of Na<sup>+</sup>/Ca<sup>2+</sup> exchange in endothelin-1-induced increases in Ca<sup>2+</sup> transient and contractility in rabbit ventricular myocytes: pharmacological analysis with KB-R7943. *Br J Pharmacol.* 1999;126:1785-95.
26. He JQ, Pi Y, Walker JW, Kamp TJ. Endothelin-1 and photoreleased diacylglycerol increase L-type Ca<sup>2+</sup> current by activation of protein kinase C in rat ventricular myocytes. *J Physiol.* 2000;52(4 Pt 3):807-20.
27. Kramer BK, Smith TW, Kelly RA. Endothelin and increased contractility in adult rat ventricular myocytes. Role of intracellular alkalosis induced by activation of the protein kinase C-dependent Na<sup>(+)</sup>-H<sup>+</sup> exchanger. *Circ Res.* 1991;68:269-79.
28. Zhang YH, James AF, Hancox JC. Regulation by endothelin-1 of Na<sup>+</sup>-Ca<sup>2+</sup> exchange current (I<sub>(NaCa)</sub>) from guinea-pig isolated ventricular myocytes. *Cell Calcium.* 2001;30:351-60.
29. Imahashi K, Pott C, Goldhaber JJ, Steenbergen C, Philipson KD, Murphy E. Cardiac-specific ablation of the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger confers protection against ischemia/reperfusion injury. *Circ Res.* 2005;97:916-21.
30. Brunner F, Opie LH. Role of endothelin-A receptors in ischemic contracture and reperfusion injury. *Circulation.* 1998;97:391-8.
31. Morawietz H, Szibor M, Goettsch W, Bartling B, Barton M, Shaw S, et al. Unloading of the left ventricle by ventricular assist device normalizes increased expression of endothelin ET(A) receptors but not endothelin-converting enzyme-1 in patients with end-stage heart failure. *Circulation.* 2000;102(19 Suppl 3):III188-93.
32. Watanabe T, Suzuki N, Shimamoto N, Fujino M, Imada A. Contribution of endogenous endothelin to the extension of myocardial infarct size in rats. *Circ Res.* 1991;69:370-7.
33. Tonnessen T, Giaid A, Saleh D, Naess PA, Yanagisawa M, Christensen G. Increased in vivo expression and production of endothelin-1 by porcine cardiomyocytes subjected to ischemia. *Circ Res.* 1995;76:767-72.
34. Galiuto L, DeMaria AN, del Balzo U, May-Newman K, Flaim SF, Wolf PL, et al. Ischemia-reperfusion injury at the microvascular level: treatment by endothelin A-selective antagonist and evaluation by myocardial contrast echocardiography. *Circulation.* 2000;102:3111-6.
35. Meyns B, Stolinski J, Leunens V, Verbeken E, Flameng W. Left ventricular support by catheter-mounted axial flow pump reduces infarct size. *J Am Coll Cardiol.* 2003;41:1087-95.
36. Yamazaki T, Komuro I, Kudoh S, Zou Y, Shiojima I, Hiroi Y, et al. Endothelin-1 is involved in mechanical stress-induced cardiomyocyte hypertrophy. *J Biol Chem.* 1996;271:3221-8.
37. Loennechen JP, Stoylen A, Beisvag V, Wisloff U, Ellingsen O. Regional expression of endothelin-1, ANP, IGF-1, and LV wall stress in the infarcted rat heart. *Am J Physiol Heart Circ Physiol.* 2001;280:H2902-10.
38. Zhang J, Narula J. Molecular biology of myocardial recovery. *Surg Clin North Am.* 2004;84:223-42.
39. Qin Y, Vanden Hoek TL, Wojcik K, Anderson T, Li CQ, Shao ZH, et al. Caspase-dependent cytochrome c release and cell death in chick cardiomyocytes after simulated ischemia-reperfusion. *Am J Physiol Heart Circ Physiol.* 2004;286:H2280-6.
40. Kajstura J, Cheng W, Reiss K, Clark WA, Sonnenblick EH, Krajewski S, et al. Apoptotic and necrotic myocyte cell deaths are independent contributing variables of infarct size in rats. *Lab Invest.* 1996;74:86-107.
41. Gottlieb RA, Engler RL. Apoptosis in myocardial ischemia-reperfusion. *Ann N Y Acad Sci.* 1999;874:412-26.
42. Oie E, Clausen OP, Yndestad A, Groggaard HK, Attramadal H. Endothelin receptor antagonism attenuates cardiomyocyte apoptosis after induction of ischemia in rats. *Scand Cardiovasc J.* 2002;36:108-16.
43. Shichiri M, Yokokura M, Marumo F, Hirata Y. Endothelin-1 inhibits apoptosis of vascular smooth muscle cells induced by nitric oxide and serum deprivation via MAP kinase pathway. *Arterioscler Thromb Vasc Biol.* 2000;20:989-97.
44. Gourine AV, Molosh AI, Poputnikov D, Bulhak A, Sjoquist PO, Pernow J. Endothelin-1 exerts a preconditioning-like cardioprotective effect against ischaemia/reperfusion injury via the ET(A) receptor and the mitochondrial K(ATP) channel in the rat in vivo. *Br J Pharmacol.* 2005;144:331-7.
45. Orrenius S, Zhivotovsky B, Nicotera P. Regulation of cell death: the calcium-apoptosis link. *Nat Rev Mol Cell Biol.* 2003;4:552-65.
46. Miyamae M, Camacho SA, Weiner MW, Figueredo VM. Attenuation of postischemic reperfusion injury is related to prevention of [Ca<sup>2+</sup>]<sub>i</sub>m overload in rat hearts. *Am J Physiol.* 1996;271:H2145-53.
47. Chen SJ, Bradley ME, Lee TC. Chemical hypoxia triggers apoptosis of cultured neonatal rat cardiac myocytes: modulation by calcium-regulated proteases and protein kinases. *Mol Cell Biochem.* 1998;178:141-9.
48. Schaper W, Flameng W, De Brabander M. Comparative aspects of coronary collateral circulation. *Adv Exp Med Biol.* 1972;22:267-76.
49. Maxwell MP, Hears DJ, Yellon DM. Species variation in the coronary collateral circulation during regional myocardial ischaemia: a critical determinant of the rate of evolution and extent of myocardial infarction. *Cardiovasc Res.* 1987;21:737-46.